

Pilot testing of a sampling methodology for assessing seed attachment propensity and transport rate in a soil matrix carried on boot soles and bike tires

Nigel Hardiman^{1, 2*}, Kristina Charlotte Dietz³, Ian Bride¹, Louis Passfield³

¹ School of Anthropology and Conservation, University of Kent, Canterbury, Kent CT2 7NZ, U.K.

² Current address: Lincoln International Business School, University of Lincoln, Brayford Pool, Lincoln LN6 7TS, U.K.

³ School of Sport and Exercise Science, University of Kent, Canterbury, Kent CT2 7NZ, U.K.

*Corresponding author: Dr. Nigel Hardiman

Lincoln International Business School, University of Lincoln

Brayford Pool, Lincoln LN6 7TS, U.K.

Phone: +44 1542 283 5527

Email: nhardiman@lincoln.ac.uk

ABSTRACT

Dietz

Keywords: weeds; seed attachment; human-mediated dispersal; tourism impacts

1. Introduction

Invasive alien species of plants (weeds), together with animals, fungi and microbes are widely recognised as posing a major threat to global biodiversity, second only to habitat destruction in their impact (Randall, 1996; Vilà *et al*, 2011; Wittenberg and Cock, 2001; World Conservation Union [IUCN], 2000). Weeds have been shown to cause billions of dollars of annual economic loss in agriculture and forestry (Pimentel *et al*, 2001; Pimentel, 2002; Williams *et al*, 2010). They have also been shown to alter ecological processes, degrade ecosystem services and disrupt ecological integrity (DiTomaso, 2000; Mack and D'Antonio, 1998; Pejchar and Mooney, 2009; Pimentel, 2002; Williams *et al*, 2010). Dispersal of weeds can occur via a variety of diaspores, including as adult individuals, ramets, bulbs or seeds, and can be mediated both by natural vectors, e.g., wind, rain, flowing water, animals, by humans or a combination of these (Nathan, 2006; Ridley, 1930, Wichmann *et al*, 2009). Studies have shown that dispersal of even small numbers of seeds, especially over large distances, can cause disproportionately large changes in ecological patterns (Cain, Milligan and Strand, 2000; Higgins, Nathan and Cain, 2003; Nathan, 2006).

One human activity with high potential for unintentional dispersal of weed seeds is tourism (including recreation). People today, especially in economically developed countries, have increasing time for leisure (Molitor, 2000) and international tourism has demonstrated rapid and almost continual growth in recent decades, with over 1 billion international tourists recorded in 2012 (UNWTO, 2013). Risk of human-mediated dispersal of seeds by recreation may be especially important in protected natural areas, where it may be one of only a few human activities allowed (Newsome, Moore and Dowling, 2002; Worboys, DeLacy and Lockwood, 2005) and where introduced seeds may develop into invasive environmental weeds. Research has shown an association between weed presence and tourism infrastructure in natural areas, especially adjoining roads and tracks (Pickering, Bear and Hill, 2007; Potito and Beatty, 2005, Spellerberg, 1998) and increasing weed diversity with increasing tourist visitation (Usher, 1988).

A small but growing number of studies have shown capacity for unintentional human-mediated dispersal of seeds by tourists, either attaching directly to hikers' clothing or equipment, embedded in soil picked up by vehicles, or animal dung/feed (for comprehensive reviews see Pickering and Mount, 2010; Ansong and Pickering, 2013 and 2014). The number of seeds dispersed by such vectors can be large (e.g., ~1300 on a walker's socks after only a five minute hike through roadside vegetation: Mount and Pickering, 2009) and of high species richness (e.g., > 750 species collected from various tourism-related vectors: Pickering and Mount, 2010), of which a high proportion have typically been subsequently identified as national or international invasive species (Mount and Pickering, 2009).

Despite such demonstrated potential, controlled experiments to quantify propensity for seed attachment and/or dispersal by people while hiking, either attaching directly to clothing or embedded in a soil matrix carried on boot soles, are scarce. We found only two studies that experimentally tested direct seed attachment rates on human skin/clothing (boots, socks, laces & trousers: Falinski, 1972; boots, socks, laces, trousers and bare legs: Mount and Pickering, 2009) and only a single study of seed attachment in a soil matrix carried on boot soles: Wichmann et al, 2009). We also found only four studies that experimentally tested dispersal of seeds attaching directly to clothing (trousers and shirts: Bullock and Primack, 1977; boots, socks, outer clothing and personal luggage: Lee and Chown, 2009; trousers and socks: Ansong, Pickering and Arthur, 2015; Pickering, Mount, Wichmann and Bullock, 2011) and a single study of seed dispersal via a soil matrix on boot soles (Wichmann et al, 2009). Even within the few aforementioned experimental studies on seed attachment on boots, relatively few factors affecting attachment rates appear to have been tested, i.e. distance walked (Falinski, 1972), trousered vs bare leg (Mount and Pickering, 2009) and seed species, individual walkers and boot types (Wichmann et al, 2009). Research on the effects of other potentially important factors, for example seed size, mass and morphology, soil type and condition (e.g., wet vs dry), appears to be scarce.

Alongside hiking, another recreation activity with high potential for weed seed introduction and/or dispersal is off-road cycling ('biking') (Pickering, Hill, Newsome and Leung, 2010). Biking is increasingly popular globally in backcountry/wilderness protected areas such as national parks (Burgin and Hardiman, 2012; Hardiman and Burgin, 2013) and in open access peri-urban natural areas (Chiu and Kriwoken, 2003) and its growth has led to increasing user group pressure for greater access to natural areas. Although a small number of experimental studies have attempted to measure biking's absolute and relative potential (e.g. vs hiking) for direct environmental degradation of such factors as increased soil exposure, decreased vegetation cover and/or species richness (e.g., Newsome and Davies [2009]; Pickering, Rossi and Barros [2011]; Thurston and Reader [2001]), no published studies to date have experimentally tested seed attachment or dispersal propensity on mountain bike tires, either in absolute terms or relative to boot soles.

The propensity for attachment and dispersal of seeds in a soil matrix on boot soles or bike tires is likely to differ for many reasons. Some key variables include: (i) available surface area of soles vs. tires (tires larger than boots [Thurston and Reader, 2001]); (ii) ground contact pattern (boots: discrete steps and equal distance covered by each boot; tires: continuous contact and different ground contact distance covered by front and rear tires); (iii) ground contact pressure (biker higher than walker [Thurston and Reader, 2001]); (iv) different tread patterns and depth of soles/tires; (v) distance covered (bike riders typically travel faster and further than walkers for a given time/effort); (vi) soil type and; (vii) soil condition (e.g. moisture content). The number and density of seeds available for attachment, along with differences in their size, morphology, weight and surface adhesion qualities, also potentially affect their attachment and/or dispersal rate. Field testing of such multiple variables is typically time-consuming and expensive. Researchers therefore need a sampling methodology that allows control of such variables while still representing 'real world' behaviour. This study sought to fill an existing knowledge gap by testing a potential sampling methodology for experimentally testing the absolute and relative propensity for seed attachment

and transport in a soil matrix (a) on boot soles and bike tires (b) in wet or dry soil (c) over different distances travelled.

2. Methods

2.1 Procedure

We constructed a circular, prefabricated track measuring 0.75m wide with 50mm sidewalls and external radius of 2.75m and internal radius 2.0m, giving a track centre line circumference of 14.92m and surface area of 11.18m². The track was designed to simulate the width of a typical outdoor trail and allow for a normal walking and cycling movement. Testing of different track widths and circumferences showed that this was the smallest size in which a typical bike could be ridden in a 'normal' fashion (i.e. without the riders' feet or hands touching the ground or a wall for balance support).

In real world conditions, the number and/or density of seeds available for attachment and dispersal is likely to be highly variable and affected by many external factors; definition of what is a 'realistic' and 'biologically-relevant' number and/or density is therefore situation-specific. To provide a benchmark, however, we designed our seed/soil density to be comparable to that used in the experiment by Wichmann *et al.* (2009). The aims and sampling methodologies of the two experiments were very different, however. In Wichmann *et al.*'s (2009) study, the researchers' primary focus was on measuring seed dispersal rate carried in a soil matrix in boot soles over distance, and their sampling protocol aimed to maximise initial seed attachment. They used 500g (volume unspecified, probably ~ 0.5 litre) of a 'sandy silty loam' soil, oven dried at 30°C, spread evenly in a tray (400mm x 250mm; soil depth unspecified), wetted with 50ml of water using a plant mister and stirred (moisture level unspecified). A walker then placed both shoe-clad feet in the tray and took 20 steps on the spot to pick up soil. The walker then stepped into a second tray (unspecified; assumed to be of same dimensions as Tray 1) containing 100 evenly spread seeds,

either *Brassica oleracea* [wild cabbage] or *Brassica nigra* [black mustard], again taking 20 steps on the spot. Assuming Tray 1 was filled to a soil depth of 20mm and Tray 1 and Tray 2 were of equivalent dimensions, this would suggest a soil area of 100,000mm² and density of seeds 100/100,000mm² = 0.001 seeds/mm², although the actual density of seeds exposed to the boot soles was probably much higher than this: 'probably artificially high' (Wichmann et al, 2009, p. 525, 530). The number of seeds attaching was calculated by subtracting the number left in the tray from 100, yielding the pickup rate (Wichmann et al, 2009, p. 524).

We used:

(1) 240 litres of soil spread evenly on the sampling track to an approximate depth of 20mm (0.02m depth x 11.18m² area = 0.2236m³). We used a commercially-obtained loam-based soil ("J. Arthur Bower's Topsoil" TM: William Sinclair Horticulture Limited, 2008).

(2) 11,180 'seeds' (11.18m² area x 0.001 seeds/mm² = 11,180), i.e. 50 'seeds'/litre of soil (vs at least 200 seeds/litre of soil in Wichmann et al [2009]). Wichmann et al (2009) used a *Brassica*-species seed, artificially coloured to aid on-ground identification. As artificially colouring the much larger quantity of seeds we used was impractical, we used synthetic "seed beads" ('Size 11 Japanese Toho' TM: Product code 11R43F; Beads Direct, 2013), purchased in a bright blue colour. The beads were roughly spherical in shape and sampling measurements showed a mean maximum diameter 2.1mm (SE = 0.07mm) and mean minimum diameter 1.6mm (SE = 0.09mm), making them comparable in size and shape to the *Brassica* spp. employed by Wichmann et al (2009). The beads were sprinkled evenly over the soil surface and mixed in by light raking before each sampling replicate.

The sampling track was set up indoors on the University of Kent's Canterbury campus and sampling was undertaken on the 4th, 6th and 7th September, 2013.

2.2 Design

140 The experiment was a 2×2×2 factorial design with factors Vector (“boot” vs “bike”), Soil Condition
141 (“moist” vs “wet”) and Traversal Distance (“short” vs “long”). For operational reasons (e.g., “wet”
142 and “moist” could not be randomised), testing followed a systematic sampling order: boot, moist,
143 short; boot, moist, long; bike, moist, short; bike, moist, long; boot, wet, short; boot, wet, long; bike,
144 wet, short; bike, wet, long. The complete sequence was replicated 7 times.

145 The Vector “Boot” comprised one pair of newly-purchased general purpose wellington boots
146 (“Traditional Green PVC Wellington Boot”, British size 8, heel/sole tread depth 10mm/5mm; Briers,
147 2011). “Bike” was a “hybrid” road/off road bicycle with side-pull caliper brakes and new tires (Claud
148 Butler “Urban 2000” 18” frame with Meghna “Explorer” 700mm x 38mm tires, with a tread depth
149 2mm).

150 Soil condition (MEA, 2013) was measured at the beginning, middle and end of each testing day,
151 using a Lutron soil moisture meter PMS-714 (Lutron, undated). “Moist” soil ranged between 18.7%-
152 21.6% during testing. After completion of moist testing, water was mist sprayed incrementally and
153 evenly onto the soil from a handheld garden sprayer and “wet” soil was >50% (moisture meter
154 maximum reading) throughout testing.

155 The Traversal Distance “short” test comprised one complete circuit of the track (≈15m) and a “long”
156 test comprised 10 circuits (≈150 m). Walking circuits were standardised to 25 discrete paces/circuit
157 (both feet combined). The same team member completed all walks and rides in an anticlockwise
158 direction.

159 On completion of each designated walk/ride distance, the walker stepped/bike was lifted carefully
160 into a sorting tray measuring 2300mm x 500mm x 50mm with a bright white base. Then during a
161 timed 10 minute period all the soil and beads adhering to boots/tires were carefully brushed off. The
162 beads were found (facilitated by their bright blue colour) and counted by team members using LED
163 head torches and magnifying glasses. After counting, beads were cleaned and, together with the soil

from the sorting tray, sprinkled evenly back around the track and the soil was raked over before commencing the next test.

2.3 Analyses

As the outcome variable, the number of beads attaching, is a non-negative count, data were analysed using (i) one-way ANOVA for testing bead attachment rate between left vs right boot soles and front vs rear bike tires; and (ii) count models (Hilbe, 2011; Ridout, Demétrio and Hinde, 1998). for testing main and interaction effects of the three factors: Vector (Boots; Tires), Soil Condition (Moist; Wet) and Traversal Distance (Short; Long); replicate number was also included in the analysis as a blocking factor, but was not significant. Poisson and negative binomial count models were considered. For several of the 8 treatment combinations, variation between replicate counts was much greater than would be expected if counts followed a Poisson distribution. Due to this over-dispersion, a negative binomial model was used for analyses of the three factors. Analyses were conducted in R, version 3.1.1 (R Core team, 2014). Results were accepted as significant at or below the 5 % probability level.

3. Results

Beads were only recorded attaching to boots and tires along with soil; no “bead-only” attachment was recorded under any sampling parameter combination. We observed that boots predominantly tended to pick up soil and beads in the heel treads, with soil tightly compacted and requiring beads to be physically extracted by the researchers, with very few beads (estimated <5%) attaching to the remainder of the soles. One-way ANOVA testing revealed no significant difference in bead attachment quantity or % rate between left and right boots for all parameter combinations ($F_{1,54} = 1.49$, $P = 0.23$). In contrast, bike tires showed a significant difference ($F_{1,54} = 15.30$, $P < 0.0003$) in bead attachment quantity and % attachment rate between front and rear tires, with attachment on the front tyre at least an order of magnitude higher than the rear for all sampling parameter

combinations except “short traversal, moist soil” (zero bead attachment recorded on both tires for all replicates, see Table 1 and 2).

The negative binomial model provided adequate fit for the data; that is predicted seed-counts did not differ significantly from the observed data, $\chi^2(49) = 62.49$, $p < 0.093$. Observed bead counts and attachment rates are therefore reported here (Table 1 and 2). Model-parameters, fit-indices and selection-criteria for the negative binomial model are reported, together with significance values for each effect, in Table 3. The model’s intercept represents an arbitrarily chosen baseline for comparison, in this case the bike/long/moist condition. The log-coefficient for the intercept represents the estimated number of seeds in that condition once exponentiated, so $\exp(0.81) = 2.25$ seeds in the bike/long/moist condition. As previously mentioned, model-estimated and actual number of seeds (2.9) did not significantly differ and, therefore, actual seed numbers are reported in Table 1. Condition effects in the model are calculated by adding relevant coefficients for main- and interaction-effects to the baseline before exponentiation. For example, to calculate the estimated number of seeds in the boot/long/wet condition, we added estimates for the Intercept, Vector, Soil Condition, and Vector x Soil Condition: $\exp(0.81+1.70+2.83+(-2.23)) = 22.42$ seeds, actual seed number = 24.7. Note that significant main effects should not be interpreted in the negative binomial model in the presence of significant interactions as they may be misleading. Condition analyses showed that, whilst there were significant effects of each of the three experimental factors (Vector, Soil Condition and Traversal Distance), all but one (Soil Condition x Traversal Distance) of the interactions between these factors were also statistically significant (Table 3). Owing to the complexity of these results and to avoid extensive statistical copy, results are summarised in the following plain text. Consistently more beads attached over the long traversal distance than over the short traversal distance; however the ratio of short to long was variable. More beads attached under wet conditions than under moist conditions, although again the ratio of wet to moist was variable. Generally, more beads attached to boots than to bike tires under the same conditions, but again the ratio was variable and this pattern reversed under the long wet conditions (Table 1). In summary,

bead attachment was higher for longer traversals and under wet soil conditions. Bead attachment was generally higher on boots than on tires, except when traversal distance was long and the soil condition was wet. Mean % attachment rate of beads from total available (11,180) was very low over all treatment combinations, ranging from 0.07% (SE = 0.02%) – 0.22% (SE = 0.03%) for boots and 0.00% (SE = 0.00%) – 0.31% (SE = 0.04%) for tires (Table 1).

4. Discussion

Our finding that bike tires had a lower propensity than boot soles to pick up beads under all conditions tested except over 150m distance travelled in wet soil was initially surprising and counter-intuitive, given the tires' larger overall surface area than the boot soles. However, the result that the bike tires tended to pick up fewer beads than boot soles makes sense, as the tread depth of the tires was shallower (2mm) than that of the boots (sole 5mm; heel 10mm) and hence the beads/soil may not have adhered as tightly to the tires as they did to the bottom of the boot. This is supported by the observation reported during testing that beads attaching to boot soles were predominantly in the heel treads (see Results above). It may be that for shorter distances and/or dryer soils the potentially deeper and narrower tread of the boot soles meant that more beads were retained on boots, but that on a longer rider on wet soil, the greater surface area of the tyre becomes more important, allowing soil to attach over a greater area resulting in more beads attaching. Increasing the density of beads in the soil in a repeat experiment so there are fewer zeros and low numbers attaching may assist in testing this hypothesis.

It must also be remembered that beads were only picked up along with soil in our experiment. It is possible that in other circumstances, for example seeds growing on trackside vegetation and possessing traits affecting attachment on walkers'/riders' clothing, for example differing morphology, mass and infructescence height might affect attachment rate, as might walkers' and riders' relative speed of travel along such tracks.

Our study gives the first published quantification of the propensity for attachment of a seed analog on bike tires, both in absolute terms and comparative to boot soles. It provides a comparison with the very small number of controlled experiments quantifying seed attachment rate on footwear, either directly or in a soil matrix, for a measured sampling effort (e.g., compare Mount and Pickering, 2009; Wichmann et al, 2009). However, comparison of our results with previous studies must be considered relative to the respective studies' very differing sampling protocols and to several important caveats which we detail below.

Our "long" test distance ($\approx 150\text{m}$) was broadly comparable to that employed by Mount and Pickering (2009; Experiment 3) who experimentally tested seed attachment on a single pair of boots worn by a single walker over 100m ($n = 20$). Their mean seed attachment quantity on boot uppers (excluding laces) and soles combined (number attaching specifically to soles unreported) was 60.5 (SE = 26.2) (trouser leg) and 71.4 (SE = 23.6) (bare leg). Our mean observed attachment quantity and variability were substantially lower, both for boots (7.7 [SE = 1.82] – 24.7 [SE = 3.25]) and tires (0.00 [SE = 0.00] – 34.6 [SE = 4.42]) under both moist and wet soil conditions (Table 1). However, these results are not directly comparable owing to very different sampling protocols employed: in the Mount and Pickering (2009) study (i) their walker traversed Australian alpine roadside vegetation, not a walking track; (ii) they measured direct seed attachment on the boots from plants and/or loose seed on the soil surface, not in the soil matrix; (iii) soil was "relatively dry" (moisture level not reported) and no soil was collected on the boots and; (iv) seed quantity available for attachment was unknown.

A key issue in all studies attempting to quantify seed attachment rates is 'what constitutes a realistic soil seed density in natural areas?' As previously noted, our experiment employed beads of comparable size, shape and density as the seeds used by Wichmann et al (2009). Our "short" walking distance of 25 steps was also broadly comparable to their sampling protocol of 20 steps. However, as their study was primarily focused on seed dispersal distance, their sampling protocol design was

designed to maximise seed attachment and their 20 steps were repeated ‘on the spot’ in each of two small [0.4 x 0.25m²] trays containing (i) wetted soil (moisture % level not reported) and (ii) 100 seeds. They recorded high attachment rates, ranging from (Experiment 1: two seed species, one walker and boot type) 4%-93% attachment, mean 52% and 42%, variability unreported and (Experiment 2: one seed species, 10 walkers, mix of walking/Wellington boots) 26%-52% attachment, mean % and variability unreported]. The authors noted that their sampling protocol did not match the “real situation” and that their recorded attachment rates were ‘probably artificially high’ (Wichmann et al, 2009, p. 525, 530). In comparison, our observed attachment rates on boots in the short distance test, under arguably more realistic “real world” conditions, were typically two orders of magnitude lower, with means ranging 0.07% [SE = 0.02%] – 0.18% [SE = 0.03%]. Attachment rates on bike tires over the same distance were lower still, with means ranging 0.00% [SE = 0.00%] – 0.14% [SE = 0.05%] (Table 1).

5. Caveats and Conclusion

Our study suggests potential benefits of a new methodology by which researchers might cost-effectively manipulate and test the effects of different influencing factors on initial seed attachment and transport rate in a soil matrix on boot soles and bike tires, both in absolute and comparable quantities. However, our results are subject to the following important caveats.

Firstly, we were using plastic beads as an analog for seeds, not real seeds. However, seeds of different species exist in a wide range of morphologies and adhesive qualities, masses and sizes and we therefore argue that our beads can be considered as a representative analog of real seeds on all three parameters except for the small hole centring the beads. The only two previous controlled studies of direct seed attachment on boots that we found (eg Falinksi, 1972; Mount and Pickering, 2009) recorded such diversity, although neither was able to quantify attachment rate in proportion to a known available seed quantity, unlike our study. Only one other controlled study (Wichmann et al., 2009) has tested attachment propensity in a soil matrix on boot soles for pre-selected, specified

288 seed types (2: *Brassica oleracea* ssp. and *Brassica nigra*): as previously noted our beads were
289 specifically selected to be a comparable size and shape to seeds used in that study.

290 Secondly, although our use of the circular test track allowed us to simulate a realistic walking and
291 riding pattern and beads were available for attachment from on top of/within shallow surface soil,
292 similar to conditions likely to be the case in a natural environment, the methodology employed in
293 the “long” (~150m) test distance, necessitating repeatedly walking/riding the same track, meant that
294 some beads might have become attached, detached and subsequently reattached on boot soles and
295 bike tires. Although we were unable to quantify this, we regularly observed soil dropping back onto
296 the track from both boots and bike tires during circuits. This was especially marked for the bike
297 under “wet” conditions, with soil (possibly containing beads) picked up on the tires often unable to
298 pass through the caliper brake pads and subsequently ejected back onto the track. This issue was
299 probably less likely to occur for boot soles because, as previously noted, boots predominantly
300 tended to pick up soil and beads in the heel treads, with soil tightly compacted and requiring beads
301 to be physically extracted by the researchers, with very few beads (estimated <5%) attaching to the
302 remainder of the soles. In defence of the sampling methodology, however, we argue:

- 303 (i) This study is a pilot test of a potentially very flexible and cost-effective sampling methodology;
304 the possible occurrence and scale of the potential attach/detach/reattach issue would benefit
305 from further testing.
- 306 (ii) The % of beads attaching from the available bead reservoir on a ‘short’/single circuit was very
307 low overall (0.07%-0.18% boot soles; 0.00%-0.14% bike tires); this suggests that the probability
308 of the same individual beads re-attaching during multiple circuits is likely to be very low.
- 309 (iii) The ‘short’/single circuit distance test is unaffected by this potential issue and estimates of seed
310 attachment over longer distances can therefore be arrived at via simple multiplication.

311 Thirdly, time and funding limits meant that our small-scale experiment used the same, single
312 walker/rider for all tests and only 1 pair of boots and 1 bike. Boots and bike tires obviously come in

313 a very wide variety of materials, sizes and tread patterns and these may affect seed attachment rate.
314 Different walking/riding behaviour of individuals may also have an effect. Wichmann et al (2009)
315 found seed attachment rate differed significantly among different walkers and shoe type (walking
316 boots vs Wellington boots), although not among different shoe sizes.

317 For the above reasons, our results presented here are necessarily case-specific and cannot be
318 generalised more widely to define the absolute relative propensity for seed attachment and
319 transport rate in a soil matrix on boot soles and bike tires. We nevertheless suggest that the
320 methodology as trialled here shows significant promise for researchers to use it more
321 comprehensively to test the attachment rate of different seed types under a range of densities and
322 soil conditions across a variety of different compounds and sizes of boot soles and bike tires, in a
323 way that is cost-effective and that reflects real-world walker and biker behaviour.

324 **Disclosure statement**

325 No potential conflict of interest was reported by the authors.

326 **Acknowledgements**

327 The authors extend grateful thanks to Professor Martin Ridout, School of Mathematics, Statistics and
328 Actuarial Science, University of Kent, for advice on data analysis and proof reading the manuscript.
329 The authors also wish to thank editors and reviewers for their excellent suggestions which
330 strengthened this manuscript.

331 **Funding**

332 This work was supported by an internal grant from the University of Kent Faculty of Social Sciences
333 Faculty Research Fund.

Table 1. Summary of results showing absolute and comparative propensity for bead attachment (observed data) on boot soles and bike tires over seven replicated tests. Note: (i) Total number of beads attaching over all tests = 810; (ii) Total number of beads available for attaching per test = 11,180.

	Moist				Wet			
	Total # beads attaching	% of total beads attaching all tests	M # (SE) of total beads attaching	Mean % attachment of beads available (SE)	Total # beads attaching	% of total beads attaching all tests	M # (SE) of total beads attaching	Mean % attachment of beads available (SE)
Boot Short left	19				48			
Boot Short right	35				89			
Boot Short total	54	6.7	7.7 (1.82)	0.07 (0.02)	137	16.9	19.6 (3.78)	0.18 (0.03)
Boot Long left	39				88			
Boot Long right	37				85			
Boot Long total	76	9.4	10.9 (1.37)	0.10 (0.01)	173	13.2	24.7 (3.25)	0.22 (0.03)
Bike Short front	0				100			
Bike Short rear	0				7			
Bike Short total	0	0.00	0.0 (0.00)	0.00 (0.00)	107	21.4	15.3 (5.13)	0.14 (0.05)
Bike Long front	19				230			
Bike Long rear	2				12			
Bike Long total	21	2.6	2.9 (0.83)	0.03 (0.01)	242	29.9	34.6 (4.42)	0.31 (0.04)

Table 2. Summary of raw data showing actual number of beads attaching on boot soles and bike tires by treatment and replicate. Total number of beads available for attaching per test = 11,180.

Boot soles							
Left moist short	Right moist short	Left moist long	Right moist long	Left wet short	Right wet short	Left wet long	Right wet long
0	0	3	5	3	4	7	8
2	2	12	6	3	14	7	14
4	3	6	3	18	11	20	22
3	5	4	4	0	11	16	10
2	7	4	9	1	18	12	8
6	9	6	5	3	15	14	8
2	9	4	5	20	16	12	15

Bike tires							
Front moist short	Rear moist short	Front moist long	Rear moist long	Front wet short	Rear wet short	Front wet long	Rear wet long
0	0	1	0	10	5	39	0
0	0	6	0	16	0	42	0
0	0	1	0	17	1	34	1
0	0	4	0	43	0	20	3
0	0	2	0	6	0	28	6
0	0	3	2	7	0	17	0
0	0	2	0	1	1	50	2

Table 3. Negative binomial model showing results of the three-factor analysis. Reported are parameter estimates (log-coefficients and associated, robust standard errors), fit- and model selection indices (LL, AIC, BIC) and associated degrees of freedom (*df*). (** = significant at $P < .01$; *** = significant at $P < .001$).

	Log-coefficient (SE)	<i>z</i>	<i>P</i>
Intercept	0.81 (0.32)**	2.893	.004
Vector	1.70 (0.32)***	5.240	<.001
Soil Condition	2.83 (0.33)***	8.944	<.001
Traversal Distance	-1.59 (0.38)***	-4.125	<.001
Vector x Traversal Distance	0.99 (0.36)**	2.994	.003
Vector x Soil Condition	-2.23 (0.34)***	-5.994	<.001
Soil Condition x Traversal Distance	0.57 (0.31)	1.649	.099
α (dispersion parameter)	0.20		
Log-Likelihood (LL)	-165.56, <i>df</i> = 8		
Akaike information criterion (AIC)	347.11, <i>df</i> = 8		
Bayesian information criterion (BIC)	363.32, <i>df</i> = 8		
Residual deviance	62.49, <i>df</i> = 49		

References

- Ansong, M. and Pickering, C. (2013). Are weeds Hitchhiking a Ride on your Car? A Systematic Review of Seed Dispersal on Cars. *PLOS ONE*, 8(11), e80275
- Ansong, M. and Pickering, C. (2014). Weed seeds on clothing: a global review. *Journal of Environmental Management*, 144(0), 203-211.
(doi: <http://dx.doi.org/10.1016/j.jenvman.2014.05.026>)
- Ansong, M., Pickering, C. and Arthur, J.M. (2015). Modelling seed retention curves for eight weed species on clothing. *Austral Ecology*, 40(7), 765-774. (doi: 10.1111/aec.12251)
- Bacela-Spychalska, K., Grabowski, M., Rewicz, T., Konopacka, A. and Wattier, R. (2013). The 'killer shrimp' *Dikerogammarus villosus* (Crustacea, Amphipoda) invading Alpine lakes: overland transport by recreational boats and scuba-diving gear as potential entry vectors? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 23(4), 606-618.
- Balmford, A., Beresford, J., Green, J., Naidoo, R., Walpole, M. and Manica, A. (2009). A Global Perspective on Trends in nature-Based Tourism. *PLoS Biology*, 7(6), e1000144, 1-6.
- Beads Direct. (2013). Size 11 Japanese Toho seed beads product specification. <http://www.beadsdirect.co.uk/beads-japanese-toho-seed-11-aqua-matt-10g-bag-11r43f/11TR43F/>. Accessed 17 September 2013.
- Briers. (2011). Home page, Briers Ltd. <http://www.briersltd.co.uk/>. Accessed 18 September 2013.
- Buckley, R. (2004). *Environmental impacts of ecotourism*. Wallingford: CABI.
- Bullock, S.H. and Primack, R.B. (1977). Comparative experimental study of seed dispersal on animals. *Ecology*, 58(3), 681–686. (doi:10.2307/1939019)
- Burgin, S. and Hardiman, N. (2012). Is the evolving sport of mountain biking compatible with nature conservation in national parks? *Australian Zoologist*, 36(2), 201-208.
- Cain, M. L., Milligan, B. G. and Strand, A. E. (2000). Long-distance seed dispersal in plant populations. *American Journal of Botany*, 87, 1217–1227. (doi:10.2307/2656714)
- Chiu, L. and Kriwoken, L. (2003). Managing recreational mountain biking in Wellington Park, Tasmania, Australia. *Annals of Leisure Research*, 6, 339-361. (doi: 10.1080/11745398.2003.10600931)

- DiTomaso, J.M. (2000). Invasive weeds in rangelands: species, impacts and management. *Weed Science*, 48, 255-265.
(doi: [http://dx.doi.org/10.1614/0043-1745\(2000\)048\[0255:IWIRSI\]2.0.CO;2](http://dx.doi.org/10.1614/0043-1745(2000)048[0255:IWIRSI]2.0.CO;2))
- Falinski, J.B. (1972). Anthropochory in xerothermic grasslands in the light of experimental data. *Acta Societatis Botanicorum Poloniae*, 41, 357–367.
- Goeft, U., Alder, J. (2001). Sustainable mountain biking: a case study from the southwest of Western Australia. *Journal of Sustainable Tourism*, 9, 193–211. (doi:10.1080/09669580108667398)
- Hardiman, N. and Burgin, S. (2010). Recreational impacts on the fauna of Australian coastal marine ecosystems. *Journal of Environmental Management*, 91, 2096-2108. (doi:10.1016/j.jenvman.2010.06.012)
- Hardiman, N. and Burgin, S. (2013). Mountain biking: is it all downhill for the environment or is there a chance to up a gear? *International Journal of Environmental Studies*, 70(6), 976-986. (doi:10.1080/00207233.2013.848531)
- Higgins, S. I., Nathan, R. and Cain, M. L. (2003). Are long-distance dispersal events in plants usually caused by nonstandard means of dispersal? *Ecology*, 84, 1945–1956. (doi:10.1890/01-0616)
- Hilbe, J. M. (2011). *Negative Binomial Regression*. Cambridge: Cambridge University Press.
- Lee, J.E. and Chown, S.L. (2009). Breaching the dispersal barrier to invasion; quantification and management. *Ecological Applications*, 19, 1944–1959. (doi.org/10.1890/08-2157.1)
- Liddle, M.J. (1997). *Recreation Ecology*. London: Chapman and Hall.
- Lutron. (Undated). Lutron Electronic Enterprise Co. Ltd. <http://www.lutron.com.tw/>. Accessed 18 September 2013.
- Mack, M.C. and D’Antonio, C.M. (1998). Impacts of biological invasions on disturbance regimes. *Trends in Ecology and Evolution*, 13(5), 195-198. (doi.org/10.1016/S0169-5347(97)01286-X)
- MEA. (2013). Learning Centre: how do you measure soil moisture content? Measurement Engineering Australia. <http://mea.com.au/soil-plants-climate/soil-moisture-monitoring/learning-centre/how-do-you-measure-soil-moisture>. Accessed 18 September, 2013.
- Molitor, G.T.T. (2000). Five economic activities likely to dominate the New Millennium: II The leisure era. *Technological Forecasting and Social Change*, 65 (2), 239–249. (doi.org/10.1016/S0040-1625(00)00112-8)
- Mount, A. and Pickering, C.M. (2009). Testing the capacity of clothing to act as a vector for non-native seed in protected areas. *Journal of Environmental Management*, 91, 168-179.

- Nathan, R. (2006). Long-distance dispersal of plants. *Science*, 313, 786–788. (doi:10.1126/science.1124975)
- Newsome, D., Moore, S.A. and Dowling, R.K. (2002). *Natural area tourism: Ecology, impacts and management*. Clevedon: Channel View.
- Newsome, D., Smith, A., Moore, S.A. (2008). Horse riding in protected areas: a critical review and implications for research and management. *Current Issues in Tourism*, 11, 1–23. (doi:10.2167/cit336.0)
- Newsome, D. and Davies, C. (2009). A case study in estimating the area of informal trail development and associated impacts caused by mountain bike activity in John Forrest National Park, Western Australia. *Journal of Ecotourism*, 8(3), 237-253. (doi:10.1080/14724040802538308)
- Pejchar, L. and Mooney, H.A. (2009). Invasive species, ecosystem services and human well-being. *Trends in Ecology and Evolution*, 24(9), 497-504. (doi.org/10.1016/j.tree.2009.03.016)
- Pickering, C.M., Bear, R. and Hill, W. (2007). Indirect impacts of nature based tourism and recreation: association between infrastructure and exotic plants in Kosciuszko National Park. *Journal of Ecotourism*, 6, 146–157. (doi:10.2167/joe162.0)
- Pickering, C. M. and Mount, A. (2010). Do tourists disperse weed seed? A global review of unintentional human-mediated terrestrial seed dispersal on clothing, vehicles and horses. *Journal of Sustainable Tourism*, 18(2), 239-256. (doi.org/10.1080/09669580903406613)
- Pickering, C.M., Hill, W. Newsome, D. and Leung, Y-F. (2010). Comparing hiking, mountain biking and horse riding impacts on vegetation and soils in Australia and the United States of America. *Journal of Environmental Management*, 91, 551-562. (doi:10.1016/j.jenvman.2009.09.025)
- Pickering, C.M., Mount, A., Wichmann, M.C. and Bullock, J.M. (2011). Estimating human-mediated dispersal of seeds within an Australian protected area. *Biological Invasions*, 13, 1869-1880.
- Pickering, C.M., Rossi, S. and Barros, A. (2011). Assessing the impacts of mountain biking and hiking on subalpine grassland in Australia using an experimental protocol. *Journal of Environmental Management*, 92(12), 3049-3057. (doi:10.1016/j.jenvman.2011.07.016)
- Pimentel, D. (2002). *Biological invasions: economic and environmental costs of alien plant, animal and microbe species*. Boca Raton: CRC Press LLC.

- Pimentel, D., McNair, S., Janecka, J., Wightman, J., Simmonds, C., O'Connell, C., Wong, E., Russel, L., Zer, J., Aquino, T. and Tsomondo, T. (2001). Economic and environmental threats of alien plant, animal, and microbe invasions. *Agriculture, Ecosystems and Environment*, 84, 1-20. (doi.org/10.1016/S0167-8809(00)00178-X)
- Potito, A.P. and Beatty, S.W. (2005). Impacts of recreation trails on exotic and ruderal species distributions in grassland areas along the Colorado Front Range. *Environmental Management*, 36, 230–236. (doi: 10.1007/s00267-003-0177-0)
- Pursuit Sports (undated). Home page. <http://www.pursuitsports.com.au/shop/home.php>. Accessed 6 February 2015.
- R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/> Accessed 6 February 2015.
- Randall, J. (1996). Weed control for the preservation of biological diversity. *Weed Technology*, 10(2), 370-383. (<http://www.jstor.org/stable/3988073>)
- Ridout, M.S., Demétrio, C.G.B. and Hinde, J. 1998. *Models for count data with many zeros*. Proceedings of the XIX International Biometric Conference, Cape Town, South Africa, 14–18 December 1998, International Biometric Society, pp. 179–192.
- Ridley, H.N. (1930). *The dispersal of plants throughout the world*. Ashford: Reeve.
- Spellerberg, I.F. (1998). Ecological Effects of Roads and Traffic: A Literature Review. *Global Ecology and Biogeography Letters*, 7(5), 317-333. (doi: 10.1046/j.1466-822x.1998.00308.x)
- The World Conservation Union (IUCN). 2000. *IUCN guidelines for the prevention of biodiversity loss caused by alien invasive species*. Species Survival Commission, Invasive Species Specialist Group. Geneva: International Union for the Conservation of Nature.
- Thurston, E. and Reader, R.J. (2001). Impacts of Experimentally Applied Mountain Biking and Hiking on Vegetation and Soil of a Deciduous Forest. *Environmental Management*, 27(3), 397-409. DOI: 10.1007/s002670010157
- Vilà, M., Espinar, J.L., Hejda, M., Hulme, P.E., Jarošík, V., Maron, J.L., Pergl, J., Schaffner, U., Sun, Y. and Pyšek, P. (2011). Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecology Letters*, 14(7), 702-728. DOI: 10.1111/j.1461-0248.2011.01628.x

- UNWTO. 2013. *UNWTO Tourism Highlights*, 2013 Edition. United Nations World Tourism Organisation. Available from: <http://mkt.unwto.org/en/publication/unwto-tourism-highlights-2013-edition>. Accessed 20 September 2013.
- Usher, M.B. (1988). Biological invasions of nature reserves: a search for generalizations. *Biological Conservation*, 44, 119–135
- Wichmann, M.C., Alexander, M.J., Soons, M.B., Galsworthy, S., Lunne, L., Gould, R., Fairfax, C., Niggemann, M., Hails, R.S. and Bullock, J.M. (2009). Human-mediated dispersal of seeds over long distances. *Proceedings of the Royal Society B*, 276, 523-532. (doi:10.1098/rspb.2008.1131)
- William Sinclair Horticulture Limited. (2008). Product data sheet, J. Arthur Bower's Topsoil TM. http://www.william-sinclair.co.uk/files/uploads/file/Gardening/COSSH%20PDFS/Other_growing_media/Top_Soil.pdf. Accessed 17 September 2013.
- Williams, F., Eschen, R., Harris, A., Djeddour, D., Pratt, C., Shaw, R.S., Varia, S., Lamontagne-Godwin, J., Thomas, S.E. and Murphy, S.T. (2010). *The Economic Cost of Invasive Non-Native Species on Great Britain*: Wallingford: CABI.
- Wittenberg, R., and Cock, M.J. (2001). *Invasive Alien Species: A Toolkit of Best Prevention and Management Practices*. Wallingford: CABI.
- Worboys, G., DeLacy, T. and Lockwood, M. (Eds.) (2005). *Protected area management: principles and practice*, 2nd edn. Cambridge: Cambridge University Press.